arXiv:1005.2993v1 [math.NT] 17 May 2010

Congruences for Hermitian modular forms of degree 2

Toshiyuki Kikuta

May 18, 2010

Abstract

We give two congruence properties of Hermitian modular forms of degree 2 over $\mathbb{Q}(\sqrt{-1})$ and $\mathbb{Q}(\sqrt{-3})$. The one is a congruence criterion for Hermitian modular forms which is generalization of Sturm's theorem. Another is the well-definedness of the p-adic weight for Hermitian modular forms.

1 Introduction and Results

1.1 Congruence criterion

Sturm [16] studied congruence properties of elliptic modular forms. In [16], the number of coefficients which are required to check the coincidence modulo a prime of two modular forms is determined. In [1], Choi and Choie studied the analog of Sturm's theorem for Jacobi forms. Choi and Choie [2], Poor and Yuen [14] and the author [8] have independently obtained some generalizations of Sturm's theorem in the case of Siegel modular forms of degree 2. The first aim of this paper is to generalize Sturm's theorem to the case of Hermitian modular forms of degree 2.

We state our results more precisely. Let $\boldsymbol{K}$ be the imaginary quadratic number field $\mathbb{Q}(\sqrt{-1})$ or $\mathbb{Q}(\sqrt{-3})$, $\mathcal{O}_{\boldsymbol{K}}$ the ring of integers in $\boldsymbol{K}$, $d_{\boldsymbol{K}}$ the discriminant of $\boldsymbol{K}$ and $U_2(\mathcal{O}_{\boldsymbol{K}})$ the Hermitian modular group of degree 2 defined as

$$U_2(\mathcal{O}_{\boldsymbol{K}}) := \left\{ M \in M_4(\mathcal{O}_{\boldsymbol{K}}) \;\middle|\; {}^t\overline{M} J_2 M = J_2 \right\}, \quad J_2 := \begin{pmatrix} O_2 & -1_2 \\ 1_2 & O_2 \end{pmatrix}.$$

We take a character ν_k on $U_2(\mathcal{O}_{\boldsymbol{K}})$ as

$$\nu_k = \begin{cases} \det^{k/2} & \text{for } \ \boldsymbol{K} = \mathbb{Q}(\sqrt{-1}), \\ \det^{k} & \text{for } \ \boldsymbol{K} = \mathbb{Q}(\sqrt{-3}). \end{cases}$$

For a subring $R \subset \mathbb{C}$, we denote by $M_k^{(s)}(U_2(\mathcal{O}_{\boldsymbol{K}}), \nu_k)_R$ the space of symmetric Hermitian modular forms of weight k with character ν_k whose all Fourier coefficients belong to R. We define $\Lambda_2(\boldsymbol{K})$ as

$$\Lambda_2(\boldsymbol{K}) := \{ H = (h_{ij}) \in Her_2(\boldsymbol{K}) \mid h_{ii} \in \mathbb{Z}, \ \sqrt{d_{\boldsymbol{K}}} h_{ij} \in \mathcal{O}_{\boldsymbol{K}} \ \},$$

where $Her_2(\boldsymbol{K})$ is the set of all Hermitian matrices of size 2 whose all components are in $\boldsymbol{K}$. For $f \in M_k(U_2(\mathcal{O}_{\boldsymbol{K}}), \nu_k)_{\mathbb{C}}$, then we write $f = \sum_H a_f(H) q^H$ the Fourier expansion of f, where $q^H := e^{2\pi i \mathrm{tr}(HZ)}$, Z is an element of the Hermitian upper-half space of degree 2, H runs over all elements of semi-positive definite of $\Lambda_2(\boldsymbol{K})$. Let p be a prime and $\mathbb{Z}_{(p)}$ the local ring of all p-integral rational numbers. We have the following theorem.

Theorem 1.1. *Let k be an even integer, p a prime with $p \geq 5$. In the case $\boldsymbol{K} = \mathbb{Q}(\sqrt{-1})$ (resp. $\boldsymbol{K} = \mathbb{Q}(\sqrt{-3})$), if $f \in M_k^{(s)}(U_2(\mathcal{O}_{\boldsymbol{K}}), \nu_k)_{\mathbb{Z}_{(p)}}$ satisfies that $a_f\left(\begin{pmatrix} m & \alpha \\ \overline{\alpha} & n \end{pmatrix}\right) \equiv 0 \bmod p$ for all m, n, α*

with $0 \leq m \leq (5k+8)/40$, $0 \leq n \leq (15k+4)/120$, $mn - N(\alpha) \geq 0$ (resp. with $0 \leq m \leq (5k+9)/45$, $0 \leq n \leq (10k+3)/90$, $mn - N(\alpha) \geq 0$), then we have $f \equiv 0 \bmod p$.

Remark 1.2. In fact, to show the assumption of the theorem, we need only to check the case $n \leq m$.

By Theorem 1.1, we obtain immediately the followings.

Corollary 1.3. Let k be an even integer, p a prime with $p \geq 5$. Let $\boldsymbol{K} = \mathbb{Q}(\sqrt{-1})$ (resp. $\boldsymbol{K} = \mathbb{Q}(\sqrt{-3})$).
(1) If $f \in M_k^{(s)}(U_2(\mathcal{O}_{\boldsymbol{K}}), \nu_k)_{\mathbb{Q}}$ satisfies that $a_f\left(\begin{pmatrix} m & \alpha \\ \overline{\alpha} & n \end{pmatrix}\right) \in \mathbb{Z}_{(p)}$ for all m, n, α with $0 \leq m \leq (5k+8)/40$, $0 \leq n \leq (15k+4)/120$, $mn - N(\alpha) \geq 0$ (resp. with $0 \leq m \leq (5k+9)/45$, $0 \leq n \leq (10k+3)/90$, $mn - N(\alpha) \geq 0$), then we have $f \in M_k(U_2(\mathcal{O}_{\boldsymbol{K}}), \nu_k)_{\mathbb{Z}_{(p)}}$.
(2) If $f \in M_k^{(s)}(U_2(\mathcal{O}_{\boldsymbol{K}}), \nu_k)_{\mathbb{Q}}$ satisfies that $a_f\left(\begin{pmatrix} m & \alpha \\ \overline{\alpha} & n \end{pmatrix}\right) = 0$ for all m, n, α with $0 \leq m \leq (5k+8)/40$, $0 \leq n \leq (15k+4)/120$, $mn - N(\alpha) \geq 0$ (resp. with $0 \leq m \leq (5k+9)/45$, $0 \leq n \leq (10k+3)/90$, $mn - N(\alpha) \geq 0$), then we have $f = 0$.

1.2 Well-definedness of p-adic weight

Serre [15] defined the notion of p-adic modular forms and applied it to the construction of p-adic L functions. Some mathematicians have attempted to generalize the theory of Serre's p-adic modular forms to the case of several variables [3, 7, 11, 13]. In particular, Ichikawa [7] showed that the p-adic weight is well-defined for Siegel modular forms of general degree. The second aim of this paper is to show the well-definedness of the p-adic weight for Hermitian modular forms of degree 2 over $\mathbb{Q}(\sqrt{-1})$ and $\mathbb{Q}(\sqrt{-3})$.

In the situation as subsection 1.1, we have the following theorem.

Theorem 1.4. Let k and k' be even integers and p a prime with $p \geq 5$. If $f \in M_k^{(s)}(U_2(\mathcal{O}_{\boldsymbol{K}}), \nu_k)_{\mathbb{Z}_{(p)}}$ and $g \in M_{k'}^{(s)}(U_2(\mathcal{O}_{\boldsymbol{K}}), \nu_{k'})_{\mathbb{Z}_{(p)}}$ satisfy that $f \not\equiv 0 \bmod p$ and $f \equiv g \bmod p^l$, then we have $k \equiv k' \bmod (p-1)p^{l-1}$.

This theorem indicates that the p-adic weight of Hermitian modular forms is "well-defined". We shall explain what the "well-defined" means. Let p be a prime, v_p the normalized additive valuation on $\mathbb{Q}_p$ (i.e. $v_p(p) = 1$) and $g = \sum_{0 \leq H \in \Lambda_2(\boldsymbol{K})} b(H) q^H$ the formal power series such that $b(H) \in \mathbb{Q}_p$ for all H. We call g a *p-adic Hermitian modular forms of degree* 2 if there exists a sequence of Hermitian modular forms $\{f_m\}$ of weight k_m with character ν_k (which are not necessarily symmetric) such that

$$\lim_{m \to \infty} f_m = g \quad (p - \text{adically}),$$

in other words,

$$\inf\{v_p(a_{f_m}(H) - b(H)) \mid 0 \leq H \in \Lambda_2(\boldsymbol{K})\} \to \infty \ (m \to \infty).$$

For simplicity, we consider only the case $p \geq 5$. We define a group $\boldsymbol{X}$ as $\boldsymbol{X} := \varprojlim \mathbb{Z}/(p-1)p^{l-1}\mathbb{Z}$ $(= \mathbb{Z}_p \times \mathbb{Z}/(p-1)\mathbb{Z})$. We find that $\mathbb{Z} \subset \boldsymbol{X}$ by the natural imbedding. The following property follows from our theorem.

Corollary 1.5. Let $\{f_m \in M_{k_m}^{(s)}(U_2(\mathcal{O}_{\boldsymbol{K}}), \nu_{k_m})_{\mathbb{Q}}\}$ and $\{f'_m \in M_{k'_m}^{(s)}(U_2(\mathcal{O}_{\boldsymbol{K}}), \nu_{k'_m})_{\mathbb{Q}}\}$ be two sequences of symmetric Hermitian modular forms of even weight. If both $\{f_m\}$ and $\{g_m\}$ converge p-adically to a same formal power series g, then there exist limits of the sequences $\{k_m\}$ and $\{k'_m\}$ in $\boldsymbol{X}$ and we have

$$\lim_{m \to \infty} k_m = \lim_{m \to \infty} k'_m \text{ in } \boldsymbol{X}.$$

By this corollary, we can define a weight of the p-adic Hermitian modular form g satisfying $g({}^tZ) = g(Z)$, similarly to the case of elliptic modular forms. In particular, we see that the p-adic limit of Hermitian Eisenstein series of weight k_m depends only on the limiting value of k_m in $\boldsymbol{X}$.

2 Hermitian modular forms

2.1 Definition and notation

The Hermitian upper half-space of degree n is defined by

$$\mathbb{H}_n := \{ Z \in M_n(\mathbb{C}) \mid \tfrac{1}{2i}(Z - {}^t\overline{Z}) > 0 \}$$

where ${}^t\overline{Z}$ is the transposed complex conjugate of Z. The space $\mathbb{H}_n$ contains the Siegel upper half-space of degree n

$$\mathbb{S}_n := \mathbb{H}_n \cap Sym_n(\mathbb{C}).$$

Let $\boldsymbol{K}$ be an imaginary quadratic number field with discriminant $d_{\boldsymbol{K}}$ and ring of integers $\mathcal{O}_{\boldsymbol{K}}$. The Hermitian modular group $U_n(\mathcal{O}_{\boldsymbol{K}})$ acts on $\mathbb{H}_n$ by the fractional transformation

$$\mathbb{H}_n \ni Z \longmapsto M < Z > := (AZ + B)(CZ + D)^{-1}, \ M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in U_n(\mathcal{O}_{\boldsymbol{K}}).$$

The subgroup $SU_n(\mathcal{O}_{\boldsymbol{K}}) := U_n(\mathcal{O}_{\boldsymbol{K}}) \cap SL_{2n}(\boldsymbol{K})$ coincides with the full group $U_n(\mathcal{O}_{\boldsymbol{K}})$ unless $d_{\boldsymbol{K}} = -3$ or -4. Let ν be an Abelian character on $U_n(\mathcal{O}_{\boldsymbol{K}})$. We denote by $M_k(U_n(\mathcal{O}_{\boldsymbol{K}}), \nu)$ the space of Hermitian modular forms of weight k and character ν for $U_n(\mathcal{O}_{\boldsymbol{K}})$. Namely, it consists of holomorphic functions $f : \mathbb{H}_n \to \mathbb{C}$ satisfying

$$f|_k M(Z) := \det(CZ + D)^{-k} f(M < Z >) = \nu(M) \cdot f(Z),$$

for all $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in U_n(\mathcal{O}_{\boldsymbol{K}})$. The cusp form $f \in M_k(U_n(\mathcal{O}_{\boldsymbol{K}}), \nu)$ is characterized by the condition

$$\Phi(f|_k \begin{pmatrix} {}^t\overline{U} & 0 \\ 0 & U \end{pmatrix}) \equiv 0 \quad \text{for all } U \in GL_n(\boldsymbol{K})$$

where Φ is the Siegel Φ-operator. A modular form $f \in M_k(U_n(\mathcal{O}_{\boldsymbol{K}}), \nu)$ is called *symmetric* if $f({}^tZ) = f(Z)$. We denote by $M_k^{(s)}(U_n(\mathcal{O}_{\boldsymbol{K}}), \nu)$ the subspace consisting of symmetric modular forms.

2.2 Fourier expansion

If $f \in M_k(U_n(\mathcal{O}_{\boldsymbol{K}}), \nu)$ satisfies the condition

$$f(Z + B) = f(Z) \quad \text{for all } B \in Her_n(\mathcal{O}_{\boldsymbol{K}}),$$

then f has a Fourier expansion of the form

$$f(Z) = \sum_{0 \le H \in \Lambda_n(\boldsymbol{K})} a_f(H) e^{2\pi i \mathrm{tr}(HZ)}$$

where

$$\Lambda_n(\boldsymbol{K}) := \{ H = (h_{ij}) \in Her_n(\boldsymbol{K}) | h_{ii} \in \mathbb{Z}, \sqrt{d_{\boldsymbol{K}}} h_{ij} \in \mathcal{O}_{\boldsymbol{K}} \}.$$

Put $\omega := \frac{1}{2}(d_{\boldsymbol{K}} + \sqrt{d_{\boldsymbol{K}}})$ and define the matrices $\dot{Z} = (\dot{z}_{ij})$ and $\ddot{Z} = (\ddot{z}_{ij})$ by

$$\dot{Z} := \frac{\omega\,{}^tZ - \bar{\omega}Z}{\omega - \bar{\omega}}, \quad \ddot{Z} := \frac{Z - {}^tZ}{\omega - \bar{\omega}}.$$

Then the above f can be considered as a function of the $\frac{1}{2}n(n-1)$ complex variables $\ddot{z}_{ij}$ $(i < j)$ in $\ddot{Z}$ and of the $\frac{1}{2}n(n+1)$ complex variables $\dot{z}_{ij}$ $(i \leq j)$ in $\dot{Z}$. Moreover, f has period 1 for each of these variables. If we define

$$\dot{q}_{ij} := \exp(2\pi i \dot{z}_{ij}) \ (i \leq j), \quad \ddot{q}_{ij} := \exp(2\pi i \ddot{z}_{ij}) \ (i < j)$$

then

$$f = \sum a_f(H) e^{2\pi i \mathrm{tr}(HZ)} = \sum a_f(H) q^H$$

may be considered as an element of the formal power series ring

$$\mathbb{C}[\dot{q}_{ij}^{\pm 1}, \ddot{q}_{ij}^{\pm 1} (i < j)][\![\dot{q}_{11}, \ldots, \dot{q}_{nn}]\!].$$

Let R be a subring of $\mathbb{C}$. We define

$$\begin{aligned} &M_k(U_n(\mathcal{O}_{\boldsymbol{K}}), \nu)_R \\ &:= \{ f = \sum a_f(H) q^H \in M_k(U_n(\mathcal{O}_{\boldsymbol{K}}), \nu) \mid a_f(H) \in R \text{ for all } H \in \Lambda_n(\boldsymbol{K}) \} \end{aligned}$$

and

$$M_k^{(s)}(U_n(\mathcal{O}_{\boldsymbol{K}}), \nu)_R := M_k(U_n(\mathcal{O}_{\boldsymbol{K}}), \nu)_R \cap M_k^{(s)}(U_n(\mathcal{O}_{\boldsymbol{K}}), \nu).$$

So we may consider the inclusion:

$$M_k(U_n(\mathcal{O}_{\boldsymbol{K}}), \nu)_R \subset R[\dot{q}_{ij}^{\pm 1}, \ddot{q}_{ij}^{\pm 1} (i < j)][\![\dot{q}_{11}, \ldots, \dot{q}_{nn}]\!].$$

3 Siegel modular forms

In this section we introduce some results concerning Siegel modular forms which are needed in later sections.

3.1 Definition and notation

Let $M_k(\Gamma_n)$ denote the space of Siegel modular forms of weight k $(\in \mathbb{Z})$ for the Siegel modular group $\Gamma_n := Sp_n(\mathbb{Z})$ and $S_k(\Gamma_n)$ the subspace of cusp forms. Any Siegel modular form $f(Z)$ in $M_k(\Gamma_n)$ has a Fourier expansion of the form

$$f(Z) = \sum_{0 \leq T \in \Lambda_n} a_f(T) e^{2\pi i \mathrm{tr}(TZ)},$$

where

$$\Lambda_n = Sym_n^*(\mathbb{Z}) := \{ T = (t_{ij}) \in Sym_n(\mathbb{Q}) \mid t_{ii},\ 2t_{ij} \in \mathbb{Z} \}$$

(the lattice in $Sym_n(\mathbb{R})$ of half-integral, symmetric matrices).
Taking $q_{ij} := \exp(2\pi i z_{ij})$ with $Z = (z_{ij}) \in \mathbb{H}_n$, we write

$$q^T := e^{2\pi i \mathrm{tr}(TZ)} = \prod_{1 \leq i < j \leq n} q_{ij}^{2t_{ij}} \prod_{i=1}^{n} q_{ii}^{t_{ii}}.$$

Using this notation, we obtain *the generalized q-expansion*:

$$f = \sum_{0 \le T \in \Lambda_n} a_F(T)\, q^T = \sum_{t_i} \left(\sum_{t_{ij}} a_f(T) \prod_{i<j} q_{ij}^{2t_{ij}} \right) \prod_{i=1}^{n} q_{ii}^{t_{ii}}$$
$$\in \mathbb{C}[q_{ij}^{-1}, q_{ij}][\![q_{11}, \ldots, q_{nn}]\!].$$

For any subring $R \subset \mathbb{C}$, we adopt the notation,

$$M_k(\Gamma_n)_R := \{ f = \sum_{T \in \Lambda_n} a_f(T) q^T \mid a_f(T) \in R \ (\forall T \in \Lambda_n) \ \},$$
$$S_k(\Gamma_n)_R := M_k(\Gamma_n)_R \cap S_k(\Gamma_n).$$

Any element $f \in M_k(\Gamma_n)_R$ can be regarded as an element of

$$R[q_{ij}^{-1}, q_{ij}][\![q_{11}, \ldots, q_{nn}]\!].$$

3.2 Siegel modular forms of degree 2

For any Siegel modular form

$$f = \sum a_f(T) q^T \in M_k(\Gamma_n)_{\mathbb{Z}_{(p)}},$$

there exists a formal power series correspondence,

$$\widetilde{f} := \sum \widetilde{a_f(T)} q^T \in \mathbb{F}_p[q_{ij}^{-1}, q_{ij}][\![q_{11}, \ldots, q_{nn}]\!].$$

where $\widetilde{a_f(T)}$ denotes the reduction modulo p of $a_f(T)$. We define

$$\widetilde{M}_k(\Gamma_n)_p : = \{ \widetilde{f} = \sum \widetilde{a_f(T)} q^T | f \in M_k(\Gamma_n)_{\mathbb{Z}_{(p)}} \}$$
$$\subset \mathbb{F}_p[q_{ij}^{-1}, q_{ij}][\![q_{11}, \ldots, q_{nn}]\!].$$

Definition 3.1. The algebra

$$\widetilde{M}(\Gamma_n)_p := \sum_{k \in \mathbb{Z}} \widetilde{M}_k(\Gamma_n)_p \quad (\textit{resp. } \widetilde{M^{(e)}}(\Gamma_n)_p := \sum_{k \in 2\mathbb{Z}} \widetilde{M}_k(\Gamma_n)_p)$$

is called *the algebra of Siegel modular forms mod p* (resp. *the algebra of Siegel modular forms mod p of even weight*).

The structure of $\widetilde{M}(\Gamma_2)_p$ was studied by Nagaoka [12]. Here we introduce the structure theorem of $\widetilde{M^{(e)}}(\Gamma_2)_p$ for the cases $p \ge 5$.

We take the usual generators G_4, G_6, X_{10} and X_{12} of the graded ring of Siegl modular forms of even weight for Γ_2. Here, G_k $(k = 4,\, 6)$ is normalized by $a_{G_k}(O_2) = 1$ and X_k $(k = 10,\, 12)$ is normalized by $a_{X_k}\left(\begin{pmatrix} 1 & \frac{1}{2} \\ \frac{1}{2} & 1 \end{pmatrix} \right) = 1$. As in [12], it is known that the graded ring over $\mathbb{Z}_{(p)}$ of Siegl modular forms of even weight with p-integral Fourier coefficients for Γ_2 is genrated by G_4, G_6, X_{10} and X_{12}.

Theorem 3.2 (Nagaoka [12]). Assume that $p \ge 5$. There exists a Siegel modular form $F_{p-1} \in M_{p-1}(\Gamma_2)_{\mathbb{Z}_{(p)}}$ such that

$$F_{p-1} \equiv 1 \bmod p$$

and one has

$$\widetilde{M^{(e)}}(\Gamma_2) \cong \mathbb{F}_p[x_1, x_2, x_3, x_4]/(\widetilde{A} - 1),$$

where $(\widetilde{A} - 1)$ is a principal ideal generated by $\widetilde{A} - 1$ and $A \in \mathbb{Z}_{(p)}[x_1, x_2, x_3, x_4]$ is defined by

$$F_{p-1} = A(G_4, G_6, X_{10}, X_{12}).$$

The following theorem is a generalization of Sturm's theorem in the case of Siegel modular forms of degree 2.

Theorem 3.3 (Kikuta [8])**.** Let p be a prime with $p \geq 5$, k an even integer and Γ an arbitrary congruence subgroup of Γ_2. We put $t := [\Gamma_2 : \Gamma]$. If $f \in M_k(\Gamma)_{\mathbb{Z}_{(p)}}$ satisfies that $a_f\left(\begin{pmatrix} m & \frac{r}{2} \\ \frac{r}{2} & n \end{pmatrix}\right) \equiv 0$ mod p for all m, n, r with $0 \leq m \leq (kt+2)/10$, $0 \leq n \leq (3kt+1)/30$ and $mn - r^2/4 \geq 0$, then we have $f \equiv 0 \bmod p$.

Remark 3.4. (1) In general, m, n, r are not integers. If Γ is a congruence subgroup of level N, then the above $\begin{pmatrix} m & \frac{r}{2} \\ \frac{r}{2} & n \end{pmatrix}$ is an element of $\frac{1}{N}\Lambda_2$.
(2) In fact, we need only to check the case $n \leq m$.

Ichikawa [7] showed that Serre's p-adic weight is well defined in the case of Siegel modular forms. We introduce the result in the general degree case.

Theorem 3.5 (Ichikawa [7])**.** Let p be a prime with $p > n+3$ or $p \equiv 1 \bmod 4$ and $f \in M_k(\Gamma_n)_{\mathbb{Z}_{(p)}}$, $g \in M_{k'}(\Gamma_n)_{\mathbb{Z}_{(p)}}$. If $f \equiv g \bmod p^l$ and $f \not\equiv 0 \bmod p$, then we have $k \equiv k' \bmod (p-1)p^{l-1}$.

4 Hermitian modular forms of degree 2

In this section, we deal with Hermitian modular forms of degree 2. For more detail, we refer to [4, 6].

We consider the Hermitian Eisenstein series of degree 2

$$E_k(Z) := \sum_{M=\left(\begin{smallmatrix} * & * \\ C & D \end{smallmatrix}\right)} (\det M)^{\frac{k}{2}} \det(CZ+D)^{-k}, \quad Z \in \mathbb{H}_2,$$

where $k > 4$ is even and $M = \left(\begin{smallmatrix} * & * \\ C & D \end{smallmatrix}\right)$ runs over a set of representatives of $\left\{\left(\begin{smallmatrix} * & * \\ O_2 & * \end{smallmatrix}\right)\right\} \backslash U_2(\mathcal{O}_{\boldsymbol{K}})$. Then we have

$$E_k \in M_k^{(s)}(U_2(\mathcal{O}_{\boldsymbol{K}}), \det^{-\frac{k}{2}}).$$

Moreover $E_4 \in M_4^{(s)}(U_2(\mathcal{O}_{\boldsymbol{K}}), \det^{-2})$ is constructed by the Maass lift ([10]).

In the rest of this paper, we mainly deal with $\boldsymbol{K} = \mathbb{Q}(\sqrt{-1})$ or $\mathbb{Q}(\sqrt{-3})$ and

$$\nu_k = \begin{cases} \det^{k/2} & \text{for } \boldsymbol{K} = \mathbb{Q}(\sqrt{-1}), \\ \det^{k} & \text{for } \boldsymbol{K} = \mathbb{Q}(\sqrt{-3}). \end{cases}$$

We remark that ν_k is a trivial character if $\#\mathcal{O}_{\boldsymbol{K}}^\times | k$. This fact follows from that there exists a unit $\varepsilon \in \mathcal{O}_{\boldsymbol{K}}^\times$ such that $\det M = \varepsilon^2$ if $M \in U_2(\mathcal{O}_{\boldsymbol{K}})$. The graded rings over $\mathbb{C}$ of these Hermitian modular forms are studied by Dern and Krieg [6] and [4].

Proposition 4.1 (cf. [4])**.** Let $f \in M_k(U_2(\mathcal{O}_{\boldsymbol{K}}), \nu_k)$. There is the following relation between the Fourier coefficients:

$$a_{f|_{\mathbb{S}_2}}\left(\begin{pmatrix} m & \frac{r}{2} \\ \frac{r}{2} & n \end{pmatrix}\right) = \sum_{\substack{\alpha \in \mathcal{D}_{\boldsymbol{K}}^{-1} \\ \alpha + \overline{\alpha} = r \\ mn - N(\alpha) \geq 0}} a_f\left(\begin{pmatrix} m & \alpha \\ \overline{\alpha} & n \end{pmatrix}\right),$$

where $\mathcal{D}_{\boldsymbol{K}}^{-1}$ is the inverse different $\mathcal{D}_{\boldsymbol{K}}^{-1} := \frac{\mathcal{O}_{\boldsymbol{K}}}{\sqrt{d_{\boldsymbol{K}}}}$.

We define a lexicographical order for the different elements

$$H = \begin{pmatrix} m & \frac{a+bi}{2} \\ \frac{a-bi}{2} & n \end{pmatrix}, \quad H' = \begin{pmatrix} m' & \frac{a'+b'i}{2} \\ \frac{a'-b'i}{2} & n' \end{pmatrix} \in \Lambda_2(\boldsymbol{K})$$

by

$$\begin{aligned} H > H' \Longleftrightarrow & \text{ (1) } \mathrm{tr}(H) > \mathrm{tr}(H') \quad \text{or} \quad \text{(2) } \mathrm{tr}(H) = \mathrm{tr}(H'),\ m > m' \\ & \text{or} \quad \text{(3) } \mathrm{tr}(H) = \mathrm{tr}(H'),\ m = m',\ a > a' \\ & \text{or} \quad \text{(4) } \mathrm{tr}(H) = \mathrm{tr}(H'),\ m = m',\ a = a',\ b > b'. \end{aligned}$$

Let p be a prime and $f \in M_k(U_2(\mathcal{O}_{\boldsymbol{K}}), \nu_k)_{\mathbb{Z}_{(p)}}$. We define an order of f by

$$\mathrm{ord}_p(f) = \min\{H | a_f(H) \not\equiv 0 \bmod p\},$$

where the "minimum" is defined in the sense of the order defined above. If $f \equiv 0 \bmod p$, then we define $\mathrm{ord}_p(f) = \infty$.

It is not difficult to see the following property.

Lemma 4.2. One has

$$\mathrm{ord}_p(fg) = \mathrm{ord}_p(f) + \mathrm{ord}_p(g).$$

Proposition 4.3 (Kikuta-Nagaoka [9])**.** Let $\boldsymbol{K} = \mathbb{Q}(\sqrt{-1})$ or $\mathbb{Q}(\sqrt{-3})$ and p be a prime with $p \geq 5$. Then there exists a Hermitian modular form $F_{p-1} \in M_{p-1}^{(s)}(U_2(\mathcal{O}_{\boldsymbol{K}}), \nu_{p-1})_{\mathbb{Z}_{(p)}}$ such that

$$F_{p-1} \equiv 1 \bmod p.$$

Theorem 4.4 (Kikuta-Nagaoka [9])**.** Let $\boldsymbol{K} = \mathbb{Q}(\sqrt{-1})$ and assume that $p \geq 5$. There exist cusp forms χ_8, F_{10}, F_{12} of respective weight 8, 10, 12 satisfying the following.
(1) E_4, E_6, χ_8, F_{10} and F_{12} are algebraically independent and

$$\bigoplus_{0 \leq k \in 2\mathbb{Z}} M_k(U_2(\mathcal{O}_{\boldsymbol{K}}), \nu_k)_{\mathbb{Z}_{(p)}}^{sym} = \mathbb{Z}_{(p)}[E_4, E_6, \chi_8, F_{10}, F_{12}].$$

In other words, if $f \in M_k^{(s)}(U_2(\mathcal{O}_{\boldsymbol{K}}), \nu_k)_{\mathbb{Z}_{(p)}}$ (k : even), then there exists a unique polynomial $P(x_1, x_2, x_3, x_4, x_5) \in \mathbb{Z}_{(p)}[x_1, x_2, x_3, x_4, x_5]$ such that

$$f = P(E_4, E_6, \chi_8, F_{10}, F_{12}).$$

(2) $E_4|_{\mathbb{S}_2} = G_4$, $\quad E_6|_{\mathbb{S}_2} = G_6$, $\quad \chi_8|_{\mathbb{S}_2} \equiv 0$, $\quad F_{10}|_{\mathbb{S}_2} = 6X_{10}$, $\quad F_{12}|_{\mathbb{S}_2} = X_{12}$.
(3) $a_{\chi_8}\left(\begin{pmatrix} 1 & \frac{-1-i}{2} \\ \frac{-1+i}{2} & 1 \end{pmatrix}\right) = 1$, namely $\mathrm{ord}_p(\chi_8) = \begin{pmatrix} 1 & \frac{-1-i}{2} \\ \frac{-1+i}{2} & 1 \end{pmatrix}$.

Theorem 4.5 (Kikuta-Nagaoka [9])**.** Let $\boldsymbol{K} = \mathbb{Q}(\sqrt{-3})$ and assume that $p \geq 5$. There exist cusp forms F_{10}, F_{12}, χ_{18} of respective weight 10, 12, 18 satisfying the following.
(1) E_4, E_6, F_{10}, F_{12}, χ_{18} are algebraically independent and

$$\bigoplus_{0 \leq k \in 2\mathbb{Z}} M_k(U_2(\mathcal{O}_{\boldsymbol{K}}), \nu_k)_{\mathbb{Z}_{(p)}}^{sym} = \mathbb{Z}_{(p)}[E_4, E_6, F_{10}, F_{12}, \chi_{18}].$$

In other words, if $f \in M_k^{(s)}(U_2(\mathcal{O}_{\boldsymbol{K}}), \nu_k)_{\mathbb{Z}_{(p)}}$ (k : even), then there exists a unique polynomial $P(x_1, x_2, x_3, x_4, x_5) \in \mathbb{Z}_{(p)}[x_1, x_2, x_3, x_4, x_5]$ such that

$$f = P(E_4, E_6, F_{10}, F_{12}, \chi_{18}).$$

(2) $E_4|_{\mathbb{S}_2} = G_4$, $\quad E_6|_{\mathbb{S}_2} = G_6$, $\quad F_{10}|_{\mathbb{S}_2} = 2X_{10}$, $\quad F_{12}|_{\mathbb{S}_2} = 2X_{12}$, $\quad \chi_{18}|_{\mathbb{S}_2} \equiv 0$.
(3) $a_{\chi_{18}}\left(\begin{pmatrix} 2 & \frac{2i}{\sqrt{3}} \\ -\frac{2i}{\sqrt{3}} & 2 \end{pmatrix}\right) = 1$, namely $\mathrm{ord}_p(\chi_{18}) = \begin{pmatrix} 2 & * \\ * & 2 \end{pmatrix}$ for some $*$.

For explicit expressions of above generators, see [9].

5 Proofs

In this section, we shall prove our theorems. However, since the proof is similar, we prove only the case $\boldsymbol{K} = \mathbb{Q}(\sqrt{-1})$.

5.1 Proof of Theorem 1.1

By Theorem 4.4 (1), we can write f in the form

$$f = P(E_4, E_6, F_{10}, F_{12}) + \chi_8 g,$$

where P is a four variables polynomial over $\mathbb{Z}_{(p)}$ and $g \in M_{k-8}^{(s)}(U_2(\mathcal{O}_{\boldsymbol{K}}), \nu_{k-8})_{\mathbb{Z}_{(p)}}$. Restricting both sides to $\mathbb{S}_2$, we obtain $f|_{\mathbb{S}_2} = P(G_4, G_6, 6X_{10}, X_{12})$ because of Theorem 4.4 (2). Proposition 4.1 implies that $a_{f|_{\mathbb{S}_2}}\left(\begin{pmatrix} m & * \\ * & n \end{pmatrix}\right) \equiv 0 \bmod p$ for all m, n with $0 \leq m \leq (5k+8)/40$ and $0 \leq n \leq (15k+4)/120$. Note that $(k+2)/10 < (5k+8)/40$ and $(3k+1)/30 < (15k+4)/120$. By Theorem 3.3, we have $f|_{\mathbb{S}_2} = P(G_4, G_6, 6X_{10}, X_{12}) \equiv 0 \bmod p$. By Theorem 3.2, we have $\widetilde{P}(x_1, x_2, 6x_3, x_4) \in (\widetilde{A} - 1)$. Since $\widetilde{P}$ is an isobaric polynomial and $6 \in \mathbb{Z}_{(p)}^{\times}$, we obtain $\widetilde{P} = 0$. Accordingly $P(E_4, E_6, F_{10}, F_{12}) \equiv 0 \bmod p$. Hence $a_{\chi_8 g}\left(\begin{pmatrix} m & * \\ * & n \end{pmatrix}\right) \equiv 0$ for all m, n with $0 \leq m \leq (5k+8)/40$, $0 \leq n \leq (15k+4)/120$.

Now we assume that the diagonal of $\mathrm{ord}_p(g\chi_8)$ are m_0 and n_0:

$$\mathrm{ord}_p(g\chi_8) = \begin{pmatrix} m_0 & * \\ * & n_0 \end{pmatrix}.$$

Note that $m_0 > (5k+8)/40$ and $n_0 > (15k+4)/120$. By Lemma 4.2, we have

$$\mathrm{ord}_p(g) = \begin{pmatrix} m_0 - 1 & * \\ * & n_0 - 1 \end{pmatrix}.$$

Then we see that $m_0 - 1 > (5k+8)/40 - 1 = (5(k-8)+8)/40$ and $n_0 - 1 > (15k+4)/120 - 1 = (15(k-8)+4)/120$. This means that $a_g\left(\begin{pmatrix} m & * \\ * & n \end{pmatrix}\right) \equiv 0 \bmod p$ for all m, n with $0 \leq m \leq (5(k-8)+8)/40$, $0 \leq n \leq (15(k-8)+4)/120$. Using an inductive argument on the weight, we see that $g \equiv 0 \bmod p$. This completes the proof of Theorem 1.1. □

Remark 5.1. This proof depends on the congruence criterion for Siegel modular forms of degree 2. Accordingly, if it's criterion is more sharp, then our results would be more sharp.

5.2 Proof of Theorem 1.4

By Theorem 4.4 (1), we can write f and g in the forms

$$f = P_1(E_4, E_6, F_{10}, F_{12}) + \chi_8 f_1, \quad g = Q_1(E_4, E_6, F_{10}, F_{12}) + \chi_8 g_1 \tag{5.1}$$

where P_1 and Q_1 are polynomials of four variables over $\mathbb{Z}_{(p)}$ and $f_1 \in M_{k-8}^{(s)}(U_2(\mathcal{O}_{\boldsymbol{K}}), \nu_{k-8})_{\mathbb{Z}_{(p)}}$ and $g_1 \in M_{k'-8}^{(s)}(U_2(\mathcal{O})_{\boldsymbol{K}}, \nu_{k'-8})_{\mathbb{Z}_{(p)}}$. By restricting two equations in (5.1) to $\mathbb{S}_2$ and Proposition 4.1, we have

$$P_1(G_4, G_6, 6X_{10}, X_{12}) \equiv Q_1(G_4, G_6, 6X_{10}, X_{12}) \bmod p^l.$$

If $P_1(G_4, G_6, 6X_{10}, X_{12}) \not\equiv 0 \bmod p$, then we obtain $k \equiv k' \bmod (p-1)p^{l-1}$ by Theorem 3.5. Hence we take the largest number $s \leq l$ such that $P_1(G_4, G_6, 6X_{10}, X_{12}) \equiv Q_1(G_4, G_6, 6X_{10}, X_{12}) \equiv 0$

mod p^s. By Theorem 3.2, we obtain $\tilde{P}_1(x_1, x_2, 6x_3, x_4) \in (\tilde{A}-1)$. Since $\tilde{P}_1$ is an isobaric polynomial, we have $\tilde{P}_1 = 0$, namely $P_1 \equiv 0 \bmod p$ as a polynomial. We obtain also $P_1 \equiv 0 \bmod p^s$ inductively. We set

$$P_1' := \frac{1}{p^s} P_1, \quad Q_1' := \frac{1}{p^s} Q_1 \in \mathbb{Z}_{(p)}[x_1, x_2, x_3, x_4].$$

By the property of s, we have

$$\begin{aligned} &P_1'(G_4, G_6, 6X_{10}, X_{12}) \equiv Q_1'(G_4, G_6, 6X_{10}, X_{12}) \bmod p^{l-s}, \\ &P_1'(G_4, G_6, 6X_{10}, X_{12}) \not\equiv 0 \bmod p. \end{aligned}$$

By Theorem 3.5, we obtain $k \equiv k' \bmod (p-1)p^{l-s-1}$. Now we may assume that $k = k' + p^{l-s-1}(p-1)a$. Taking multiplication of F_{p-1} which is given in Proposition 4.3, we have

$$Q_1'(E_4, E_6, F_{10}, F_{12}) \equiv Q_1'(E_4, E_6, F_{10}, F_{12}) F_{p-1}^{ap^{(l-s-1)}} \bmod p^{l-s}. \tag{5.2}$$

By Theorem 4.4 (1), we can write

$$Q_1'(E_4, E_6, F_{10}, F_{12}) F_{p-1}^{ap^{(l-s-1)}} = R_1(E_4, E_6, F_{10}, F_{12}) + \chi_8 h_1,$$

where R_1 is a four variables polynomial over $\mathbb{Z}_{(p)}$ and $h \in M_{k-8}(U_2(\mathcal{O})_{\boldsymbol{K}}, \nu_{k-8})$. Note that $P_1'(E_4, E_6, F_{10}, F_{12})$, $R_1(E_4, E_6, F_{10}, F_{12}) + \chi_8 h_1 \in M_k^{(s)}(U_2(\mathcal{O}_{\boldsymbol{K}}), \nu_k)_{\mathbb{Z}_{(p)}}$. Since (5.2) and by using Theorem 4.4 (2), we have

$$P_1'(G_4, G_6, 6X_{10}, X_{12}) \equiv R_1(G_4, G_6, 6X_{10}, X_{12}) \bmod p^{l-s}.$$

Since the weights of both sides are same, by using Theorem 3.2 repeatedly, we get $P_1'(E_4, E_6, F_{10}, F_{12}) \equiv R_1(E_4, E_6, F_{10}, F_{12}) \bmod p^{l-s}$. It follows that

$$P_1(E_4, E_6, F_{10}, F_{12}) \equiv p^s R_1(E_4, E_6, F_{10}, F_{12}) \equiv Q_1(E_4, E_6, F_{10}, F_{12}) - p^s \chi_8 h_1 \bmod p^l.$$

By the assumption that $f \equiv g \bmod p^l$, we have

$$\chi_8(f_1 - h_1 p^s) \equiv \chi_8 g_1 \bmod p^l.$$

Now we substitute $f_1 - h_1 p^s$ to f_1. Since $f \not\equiv 0 \bmod p$, it must be that $\chi_8 f_1 \not\equiv 0 \bmod p$. By Lemma 4.2, we see also that $f_1 \not\equiv 0 \bmod p$ and $f_1 \equiv g_1 \bmod p^l$. We write again f_1 and g_1 in the forms

$$f_1 = P_2(E_4, E_6, F_{10}, F_{12}) + \chi_8 f_2, \quad g_1 = Q_2(E_4, E_6, F_{10}, F_{12}) + \chi_8 g_2,$$

where P_2 and Q_2 are polynomials of four variables over $\mathbb{Z}_{(p)}$, $f_2 \in M_{k-16}^{(s)}(U_2(\mathcal{O}_{\boldsymbol{K}}), \nu_{k-16})_{\mathbb{Z}_{(p)}}$ and $g_2 \in M_{k'-16}^{(s)}(U_2(\mathcal{O}_{\boldsymbol{K}}), \nu_{k'-16})_{\mathbb{Z}_{(p)}}$. By Theorem 3.5, if $P_2(G_4, G_6, 6X_{10}, X_{12}) \not\equiv 0 \bmod p$, then we obtain $k - 8 \equiv k' - 8 \bmod (p-1)p^{l-1}$, namely $k \equiv k' \bmod (p-1)p^{l-1}$. If otherwise, by continuing the above argument repeatedly, we find a natural number t such that $g_t|_{\mathbb{S}_2} \not\equiv 0 \bmod p$. In fact, it suffices to continue until t satisfying $0 \le k' - 8t < 8$. Applying Theorem 3.5 to f_t and g_t, we obtain that $k - 8t \equiv k' - 8t \bmod (p-1)p^{l-1}$, namely $k \equiv k' \bmod (p-1)p^{l-1}$. This completes the proof of Theorem 1.4. $\square$

References

[1] D. Choi, Y. Choie, On the congruences of Jacobi forms. Math. Z. 256 (2007) 287-294.

[2] D. Choi, Y. Choie, The fourier coefficients of Siegel modular forms modulo p, preprint.

[3] S. Böcherer, S. Nagaoka, On mod p properties of Siegel modular forms, Math. Ann. 338 (2007) 421-433.

[4] T. Dern, Hermitesche Modulformen zweiten Grades, Verlag Mainz, Wissenschaftsverlag, Aachen (2001)

[5] J-I. Igusa, Siegel modular forms of genus two, Amer. J. Math. 84 (1962) 175-200.

[6] T. Dern, A. Krieg, Graded rings of Hermitian modular forms of degree 2, manuscipta math. 110 (2003) 251-272.

[7] T. Ichikawa, Congruences between Siegel modular forms, Math. Ann. 342 (2008) 527-532.

[8] T. Kikuta, Sturm type theorem for Siegel modular forms of degree 2, preprint.

[9] T. Kikuta, Nagaoka, S.: Hermitian modular forms mod p, preprint.

[10] A. Krieg, The Maass spaces on the Hermitian half-space of degree 2, Math. Ann. 289 (1991) 663-681.

[11] S. Nagaoka, On p-adic Hermitian Eisenstein series, Proc. AMS. 134 (2006) 2533-2540.

[12] S. Nagaoka, Note on mod p Siegel modular forms I, II, Math. Z. 235 (2000) 227-250, ibid. 251 (2005) 821–826.

[13] T. Munemoto, S. Nagaoka, Note on p-adic Hermitian Eisenstein series, Abh. Math. Univ. Hambg 76 (2006) 247-260.

[14] C. Poor, D. Yuen, Paramodular cusp forms, preprint.

[15] J.-P. Serre, Formes modulaires et fonctions zêta p-adiques, Modular functions of one variable V, Springer Lect. Notes in Math. 350 (1973) 191-268.

[16] J. Sturm, On the congruence of modular forms, Springer Lect. Notes in Math. 1240 (1984) 275-228.

T. Kikuta
Department of Mathematics Kinki University Higashi-Osaka, 577-8502 Osaka, Japan
Tel.: +81-6-6721-2332
Fax: +81-6-6727-4301
E-mail: kikuta84@gmail.com